



MISSISSIPPI STATE DEPARTMENT OF HEALTH

**This is an official
MS Health Alert Network (HAN) Alert**

MESSAGE ID: MSHAN-20210610-00522-ALT (Health Alert)
RECIPIENTS: All Physicians, Hospitals, ERs, ICPs, NPs, and
Healthcare Providers – Statewide
Thursday, June 10, 2021
SUBJECT: Increased Respiratory Syncytial Virus Activity in
the Southern US

Dear Colleagues,

- The Centers for Disease Control and Prevention (CDC) has issued the below Health Advisory notifying clinicians of increased interseasonal respiratory syncytial virus (RSV) activity, particularly in the Southern US.
- In the US, RSV typically is a fall and winter illness. Beginning in April 2020, RSV activity rapidly declined and remained low, like many other respiratory illnesses including influenza. **Those declines in activity were likely due to the many public health measures in place to prevent COVID-19 transmission, such as universal mask use and physical distancing.**
- Since late March 2021, CDC has seen increases in RSV activity around the US, primarily in the South, including HHS Region 4 which contains Mississippi. See National and Regional RSV data available at the [National Respiratory and Enteric Virus Surveillance System](#)
- Due to reduced circulation of RSV in the previous fall and winter, older infants and toddlers might now be at increased risk of severe RSV-associated illness since they have likely not had typical levels of exposure to RSV during the past 15 months.
- The Mississippi State Department of Health has received reports of several RSV cases in childcare facilities, highlighting the potential for increased activity and transmission in these settings.
- Other settings of concern include long-term care facilities.

In Mississippi, Physicians and Providers as asked to:

- Consider RSV and other respiratory pathogens among patients presenting with acute respiratory illness who test negative for SARS-CoV-2, especially children and those in congregate settings. Real-time reverse transcription-polymerase chain reaction (RT-PCR) is the preferred method for testing for respiratory viruses.
- Remind patients, especially healthcare personnel, childcare providers, and staff of long-term care facilities to not go to work while acutely ill, even if they have a negative COVID-19 test.



- Report suspected clusters or outbreaks of RSV to the Mississippi State Department of Health Office of Epidemiology at 601-576-7725.
- Please see the full CDC Health Advisory for additional details, guidance, and resources.

Regards,

Paul Byers, MD
State Epidemiologist

This is an official **CDC HEALTH ADVISORY**

Distributed via the CDC Health Alert Network
June 10, 2021, 1 :30 EST PM
CDCHAN-00443

Increased Interseasonal Respiratory Syncytial Virus (RSV) Activity in Parts of the Southern United States

Summary

The Centers for Disease Control and Prevention (CDC) is issuing this health advisory to notify clinicians and caregivers about increased interseasonal respiratory syncytial virus (RSV) activity across parts of the Southern United States. Due to this increased activity, CDC encourages broader testing for RSV among patients presenting with acute respiratory illness who test negative for SARS-CoV-2, the virus that causes COVID-19. RSV can be associated with severe disease in young children and older adults. This health advisory also serves as a reminder to healthcare personnel, childcare providers, and staff of long-term care facilities to avoid reporting to work while acutely ill – even if they test negative for SARS-CoV-2.

Background

RSV is an RNA virus of the genus *Orthopneumovirus*, family *Pneumoviridae*, primarily spread via respiratory droplets when a person coughs or sneezes, and through direct contact with a contaminated surface. RSV is the most common cause of bronchiolitis and pneumonia in children under one year of age in the United States. Infants, young children, and older adults with chronic medical conditions are at risk of severe disease from RSV infection. Each year in the United States, RSV leads to on average approximately 58,000 hospitalizations¹ with 100-500 deaths among children younger than 5 years old² and 177,000 hospitalizations with 14,000 deaths among adults aged 65 years or older.³

In the United States, RSV infections occur primarily during the fall and winter cold and flu season. In April 2020, RSV activity decreased rapidly, likely due to the adoption of public health measures to reduce the spread of COVID-19.⁴ Compared with previous years, RSV activity remained relatively low from May 2020 to March 2021. However, since late March, CDC has observed an increase in RSV detections reported to the [National Respiratory and Enteric Virus Surveillance System](#) (NREVSS), a nationwide passive, laboratory-based surveillance network. CDC noted increases in laboratory detections and in the percentages of positive detections for both antigen and PCR testing in parts of HHS Region 4 (Alabama,

570 East Woodrow Wilson • Post Office Box 1700 • Jackson, MS 39215-1700
601-576-8090 • 1-866-HLTHY4U • www.HealthyMS.com



Florida, Georgia, Kentucky, Mississippi, North Carolina, South Carolina, and Tennessee) and Region 6 (Arkansas, Louisiana, New Mexico, Oklahoma, and Texas). Due to limited testing outside of the typical RSV season, data are limited in some jurisdictions and may be incomplete for the most recent weeks. Since this elevated interseasonal activity is a deviation in the typical circulation patterns for RSV, at this time it is not possible to anticipate the likely spread, peak, or duration of activity with any certainty. Health officials also identified increased interseasonal RSV circulation in parts of Australia during late 2020 and in South Africa in early 2021. Still, RSV did not reach seasonal peak levels in most regions or result in widespread circulation.⁵⁻⁷

Due to reduced circulation of RSV during the winter months of 2020–2021, older infants and toddlers might now be at increased risk of severe RSV-associated illness since they have likely not had typical levels of exposure to RSV during the past 15 months. In infants younger than six months, RSV infection may result in symptoms of irritability, poor feeding, lethargy, and/or apnea with or without fever. In older infants and young children, rhinorrhea and decreased appetite may appear one to three days before cough, often followed by sneezing, fever, and sometimes wheezing. Symptoms in adults are typically consistent with upper respiratory tract infections, including rhinorrhea, pharyngitis, cough, headache, fatigue, and fever. There is no specific treatment for RSV infection other than [symptom management](#).

Recommendations

1. Clinicians and caregivers should be aware of the typical clinical presentation of RSV for different age groups.
2. Clinicians should consider testing patients with a negative SARS-CoV-2 test and acute respiratory illness or the age-specific symptoms presented above for non-SARS-CoV-2 respiratory pathogens, such as RSV. Real-time reverse transcription-polymerase chain reaction (rRT-PCR) is the preferred method for testing for respiratory viruses.
3. Clinicians should report laboratory-confirmed RSV cases and suspected clusters of severe respiratory illness to local and state health departments according to their routine reporting requirements.
4. Healthcare personnel, childcare providers, and staff of long-term care facilities should avoid reporting to work while acutely ill – even if they test negative for SARS-CoV-2.
5. Clinicians can review weekly updates to the [NREVSS website](#) and refer to surveillance data collected by local hospitals and health departments for information on RSV circulation trends in their area.

For More Information

- [CDC – RSV Information for Healthcare Providers](#)
- [CDC – RSV National Trends - NREVSS](#)
- [CDC – RSV Trends and Surveillance](#)
- [CDC – RSV Symptoms and Care](#)

References

¹[Rha B, Curns AT, Lively JY, Campbell AP, Englund JA, Boom JA, Azimi PH, Weinberg GA, Staat MA, Selvarangan R, Halasa NB, McNeal MM, Klein EJ, Harrison CJ, Williams JV, Szilagyi PG, Singer MN, Sahni LC, Figueroa-Downing D, McDaniel D, Prill MM, Whitaker BL, Stewart LS, Schuster JE, Pahud BA, Weddle G, Avadhanula V, Munoz FM, Piedra PA, Payne DC, Langley G, Gerber SI. Respiratory Syncytial Virus-Associated Hospitalizations Among Young Children: 2015-2016. *Pediatrics*. 2020 Jul;146\(1\):e20193611. doi: 10.1542/peds.2019-3611. Epub 2020 Jun 16. PMID: 32546583.](#)

²[Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, Fukuda K. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA*. 2003 Jan 8;289\(2\):179-86. doi: 10.1001/jama.289.2.179. PMID: 12517228.](#)



MISSISSIPPI STATE DEPARTMENT OF HEALTH

³[Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. N Engl J Med. 2005 Apr 28;352\(17\):1749-59. doi: 10.1056/NEJMoa043951. PMID: 15858184.](#)

⁴[Haddadin Z, Schuster JE, Spieker AJ, Rahman H, Blozinski A, Stewart L, Campbell AP, Lively JY, Michaels MG, Williams JV, Boom JA, Sahni LC, Staat M, McNeal M, Selvarangan R, Harrison CJ, Weinberg GA, Szilagyi PG, Englund JA, Klein EJ, Curns AT, Rha B, Langley GE, Hall AJ, Patel MM, Halasa NB. Acute Respiratory Illnesses in Children in the SARS-CoV-2 Pandemic: Prospective Multicenter Study. Pediatrics. 2021 May 13:e2021051462. doi: 10.1542/peds.2021-051462. Epub ahead of print. PMID: 33986150.](#)

⁵[Foley DA, Yeoh DK, Minney-Smith CA, Martin AC, Mace AO, Sikazwe CT, Le H, Levy A, Moore HC, Blyth CC. The Interseasonal Resurgence of Respiratory Syncytial Virus in Australian Children Following the Reduction of Coronavirus Disease 2019-Related Public Health Measures. Clin Infect Dis. 2021 Feb 17:ciaa1906. doi: 10.1093/cid/ciaa1906. Epub ahead of print. PMID: 33594407; PMCID: PMC7929151.](#)

⁶[Virus Watch](#), Week Ending 30 May 2021. Government of Western Australia, Department of Health. Accessed 6/7/2021.

⁷[Weekly Respiratory Pathogens Surveillance Report](#), South Africa, Week 21, 2021. National Institute for Communicable Diseases, Division of the National Health Laboratory Service. Accessed 6/7/21.

The Centers for Disease Control and Prevention (CDC) protects people's health and safety by preventing and controlling diseases and injuries; enhances health decisions by providing credible information on critical health issues; and promotes healthy living through strong partnerships with local, national, and international organizations.



Alerting Message Specification Settings

Originating Agency:	Mississippi State Department of Health
Alerting Program:	MS Health Alert Network (MS HAN)
Message Identifier:	MSHAN-20210610-00522-ALT
Program (HAN) Type:	Health Alert
Status (Type):	Actual ()
Message Type:	Alert
Reference:	MSHAN-00522
Severity:	Unknown
Acknowledgement:	No
Sensitive:	Not Sensitive
Message Expiration:	Undetermined
Urgency:	Undetermined
Delivery Time:	600 minutes

Definition of Alerting Vocabulary and Message Specification Settings

Originating Agency:	A unique identifier for the agency originating the alert.
Alerting Program:	The program sending the alert or engaging in alerts and communications using PHIN Communication and Alerting (PCA) as a vehicle for their delivery.
Message Identifier:	A unique alert identifier that is generated upon alert activation (MSHAN-yyymmdd-hhmm-TTT (ALT=Health Alert , ADV=Health Advisory , UPD=Health Update , MSG/INFO=Message/Info Service)).
Program (HAN) Type:	Categories of Health Alert Messages.
Health Alert:	Conveys the highest level of importance; warrants immediate action or attention.
Health Advisory:	Provides important information for a specific incident or situation; may not require immediate action.
Health Update:	Provides updated information regarding an incident or situation; unlikely to require immediate action.
Health Info Service:	Provides Message / Notification of general public health information; unlikely to require immediate action.
Status (Type):	
Actual:	Communication or alert refers to a live event
Exercise:	Designated recipients must respond to the communication or alert
Test:	Communication or alert is related to a technical, system test and should be disregarded



Message Type:

Alert: Indicates an original Alert
Update: Indicates prior alert has been Updated and/or superseded
Cancel: Indicates prior alert has been cancelled
Error: Indicates prior alert has been retracted

Reference: For a communication or alert with a Message Type of “Update” or “Cancel”, this attribute contains the unique Message Identifier of the original communication or alert being updated or cancelled. “n/a” = Not Applicable.

Severity:

Extreme: Extraordinary threat to life or property
Severe: Significant threat to life or property
Moderate: Possible threat to life or property
Minor: Minimal threat to life or property
Unknown: Unknown threat to life or property

Acknowledgement: Indicates whether an acknowledgement on the part of the recipient is required to confirm that the alert was received, and the timeframe in which a response is required (Yes or No).

Sensitive:

Sensitive: Indicates the alert contains sensitive content
Not Sensitive: Indicates non-sensitive content

Message Expiration: Undetermined.

Urgency: Undetermined. Responsive action should be taken immediately.

Delivery Time: Indicates the timeframe for delivery of the alert (15, 60, 1440, 4320 minutes (.25, 1, 24, 72 hours)).